



International Journal Of Scientific And University Research Publication

ISSN No **2364/2018**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



BACTERIAL AND FUNGAL STUDY OF 100 CASES OF CHRONIC SUPPURATIVE OTITIS MEDIA

Dr.R.Usha Rani || Senior Resident

Department of Microbiology

Rangaraya Medical College

Kakinada.

Objective – To study the bacterial and fungal organisms present in chronic suppurative otitis media and study the antibiotic sensitivity pattern of the bacterial

isolates. Methodology – A total number of 100 patients with clinical diagnosis of CSOM, attending the ENT OP comprised the study group. Three pus swabs from each patient discharging ear were collected. One pus swab is used for smear preparation, second swab for bacterial cultures and third swab for fungal cultures. Bacterial isolates identified and antibiotic susceptibility testing is done. Fungal isolates were identified by conventional methods. The findings were analysed. Results – Out of 100 patients, 53% were females and 47% were males. In this 40% belonged to 0-15 yrs age group. From 100 bacterial cultures 95 bacterial strains were isolated. In this *Pseudomonas aeruginosa* (36), *Klebsiella* (20), *Staphylococcus aureus* (10) followed by other organisms. Different fungal organisms isolated from 100 cultures were *Candida albicans* (12), *Aspergillus flavus* (5), *Aspergillus niger* (4). Study of antibiotic susceptibility of 95 bacterial isolates revealed that Imipenem is most effective without resistance followed by Amikacin (6.3% resistance), Gentamicin (7.3%), Ciprofloxacin (10.5%) followed by other antibiotics. Conclusion – The present study stresses the need for careful isolation of both bacterial and fungal organisms to establish the role of fungi in CSOM pathology and to prevent administration of unwanted antibiotics.

**Bacterial ,Fungal ,Study ,Chronic
Suppurative, Otitis Media.**

مقدمة

Blood agar and MacConkey agar. The bacterial strains isolated were identified according to standard procedures given in Mackie MacCartney 14th edition². Bacterial isolates were subjected to -antibiotic susceptibility test using disc diffusion method by Kirby Bauer. Fungal growth on SDA were identified by standard identification methods³.

STLUSER

Out of 100 CSOM patients investigated in the present study includes various age groups ranging from 3-70 years. In these 100 cases 47 were males and 53 were females. In this 40 cases belonged to 0-15 years age group. On culture of 100 patients swabs 95 bacterial strains and 21 fungal strains were isolated. 68 patients swabs showed growth of one bacterial organism, 4 patients swabs shows mixture of two bacterial strains, 19 patients swabs showed one bacterial and one fungal strain, 2 patients swabs shown only fungal growth and 7 patients swabs showed no growth of any organism. Out of 95 bacterial organisms isolated from cultures, *Pseudomonas aeruginosa* 36 (37.89%), *Klebsiella pneumoniae* 20 (21.05%), *Staphylococcus aureus* 10 (10.52%), *Coagulase negative Staphylococci* 9 (9.47%), *Escherichia coli* 7 (7.35%), *Proteus* 6 (6.31%), *Acinetobacter* 3 (3.15%), *Providencia* 2 (2.1%) and *Morganella morganii* 2 (2.1%). Out of 21 fungal organisms isolated from cultures, *Candida albicans* 12 (57.14%), *Aspergillus flavus* 5 (23.80%), *Aspergillus niger* 4 (19.04%) were found. Overall antibiotic sensitivity patterns of aerobic bacterial organisms from CSOM cases revealed that Imipenem (imp) was most effective antibiotic followed by Amikacin (Ak), Gentamicin (Gen), Ciprofloxacin (cip), Amoxycillin plus clavulanic acid (Amc), Cefatoxime (ctx) and Cotrimaxazole (Cot). IJSR - INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH 125 Volume : 4 | Issue : 8 | Aug 2015 • ISSN No 2277 - 8179

CSOM with its complications is one of the common diseases in the practice of otologist, paediatrician and general practitioner. CSOM is an infection of the middle ear and mastoid cavity¹. It is defined as persistent or intermittent infected discharge of more than 3 months duration through perforated or non intact tympanic membrane caused by bacteria, fungi and viruses resulting in inflammation of the mucosal lining that often results in partial or total loss of tympanic membrane and the ossicles

5. CSOM is one of the most common causes of deafness and can cause permanent perforation of tympanic membrane. World Health Organization had estimated that CSOM prevalence in India, Tanzania, Solomon islands, Guam, Australian aborigines and Greenland is highest (> 4%) and urgent attention is needed to deal with the health problem¹. In the recent years a steady increase in the incidence of gram negative bacterial infections has been observed. Many a times random antibiotic therapy prior to a defined laboratory diagnosis, leads to indiscriminate use of antibiotics, which results in emergence of resistant strains. Recent advances in the management of CSOM emphasize the necessity of bacteriological examination of ear discharge and mycological study. In the present study an earnest attempt was made to study the bacterial and fungal flora of CSOM cases.

YGOLODHOTME

A total number of 100 patients with clinical diagnosis of CSOM attending ENT OP department of Govt. General Hospital, Kakinada during February 2012 to September 2014, comprised the study group. All these are not having recent treatment with antibiotics either locally or systemically. These cases included are individuals of both sexes and all age groups. Clinical evaluation of the disease was done by presence of perforation of tympanic membrane and otorrhoea which are the two presenting symptoms of patients. The discharging pus was collected under aseptic conditions with the help of sterile swabs in triplicate by dipping into deep meatus through its drains. The pus samples from each patient was analyzed in the department of Microbiology. The material of the first swab was used for making smears and for KOH preparation for microscopic examination. The second swab was used to inoculate on two Sabouraud's Dextrose agar slopes and incubated at 37°C and another at 25°C. The third swab was used for bacteriological cultures on

– SHOWING THE BACTERIAL ABLE TResearch Paper ISOLATES FROM CSOM CASES AND THEIR RESISTANCE PATTERN TO ANTIBIOTICS

Cotrimaxazole	Amoxycillin plus clavulanic acid	Cefatoxime	Gentamicin	Imipenem	Other
17	17	5	14	36	1
47	47	14	14	36	1
0%	0%	0%	0%	0%	36%

domonas aeruginosa

NOISSISCUD

ref str

0	6	0	1	1	0	20	2.
30%			5%	5%		0	KL esi sell a pn eu m on ia e
30%	0	0	5	0	0	10	3.
			50%				St ap hy lo co cc us au re us
67%	0	0	2	0	0	09	4.
			20%				C O N S
				3			
571%	0	2	2	43%	0	07	5.
		28%	28%				E. co li
33%	1	1	0	3	0	06	6.
33%	17%	17%	50%				Pr ot e us s pe ci es
33%	1	1	0	0	0	03	7.
							A ci ne to ba ct er
	1	0	0	1	0	02	8.
	50%			50%			Pr ov id en ci a sp ec ie s
	0	0	0	0	0	02	9.
							M or ga ne lla m or ga nn ii
27	26	21	10	7	6	095	T ot al
							% of or ga
28	27	22	10	7	6	0	

- media burden of illness and management **Chronic suppurative otitis** •
- options, World Health Organisation, Geneva, Switzerland 2004.
- 1996 14th edition. Churchill Livingstone, **Mackie Mac Cartney**, •
- Edinburg Publishers.
- Text book of Medical Mycology, 3rd edition 2011. **Jagadish Chander**, •
- of Otorhinolaryngology **Scott Brown's Text book** •
- 7th ed., 1997. Butterworth Haninemann Publishers.
- , Shrestha. I, Ghosh. I, Microbiological **Shrestha. B. L., Amartyar. C. M** •
- profile of Chronic Suppurative Otitis Media Nepalese Journal of ENT, •
- Head & Neck surgery 2011, July – December, Vol. 2 Issue (2) pp. 6-7
- Sonia Seth (2011) Bacterial and fungal study of **Harvinder Kumar**, •
- 100 cases of Chronic Suppurative Otitis Media, Journal of Clinical & •
- Diagnostic Research, 2011, November (suppl-1), Vol-5(6), pp. 1224 •
- 1227.
- , Study of Bacterial flora in CSOM and **V. K. Poorey, Arathi Iyer et al** •
- its Clinical significance, Indian Journal of Otorhinolaryngology and •
- Head and neck surgery, April- June 2002, Vol – 54, No. 2, pp-91-95, •
- 2011 .
- (2000), Saudi Medical Journal **Mohammed S. Attallah** •
- Vol-21(10), pp-925.
- (2002), Microbiology of Chronic **A. H. C. Loy, A. L. Tan, P. K. S. Lu** •
- Suppurative Otitis Media, in Singapore, Singapore Medical Journal •
- 2002, Vol-43(6), pp-296-299.
- Mohammad Ismail Khan, Luqman Satt (2011), Gomal **Kamran Iqbal**, •
- Journal of Medical Sciences, Jul-Dec 2011, Vol-9, No-2. 2011. •
- and Goyal. o. p (1976). Bacteriological study in Chronic **Changani. D. L** •
- Suppurative Otitis Media, Indian Journal of •
- Otolaryngology, 28, pp-41-45.
- .(1963). Diseases of ear 1st edition, Edward Arnold **Mawson. S. R** •
- ltd. London.



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com