



International Journal Of Scientific And University Research Publication

ISSN No **301/704**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



USE OF COMPLEMENTARY AND ALTERNATIVE MEDICINE BY CANCER PATIENTS: SYSTEMATIC REVIEW

Cristiane Spadacioli || I Programa de Pós-graduação em Saúde Coletiva. Faculdade de Ciências Médicas (FCM). Universidade Estadual de Campinas (Unicamp). Campinas

**SP
Brasil.**

Interest in complementary and alternative medicine has increased, especially among oncology patients. This systematic literature review of the profile of patients who choose to use

this type of medicine, as well as their motivations, was carried out on the PubMed database. For this search, the key words used were "cancer and complementary alternative medicine" and "oncology and complementary alternative medicine", covering the period between 1995 and 2005. The selection criteria were the following: key words were present in the article title; article was written in either English, Portuguese, or Spanish; and study was performed with an adult population. From the 43 articles analyzed, it could be concluded that the use of complementary and alternative medicine is part of these patients' social identity. Moreover, its use plays an important role in the identity construction of cancer patients, helping them to make decisions related to conventional treatment.

Neoplasms, prevention & control Complementary therapies.

-However, specific debate over the socioeconomic, ethnic and gender oriented profile, as well as the patients' motivations for the use of CAM in the treatment of cancer, was not identified in the literature. The objective of the present study was to analyze the profile of people who use complementary and alternative medicine and their motivations, based on biomedical literature review of this theme.

METHODS

A literature review of this theme was performed on the PubMed of the National Library of Medicine for the ten-year period between 1995 and 2005. The key words used were: "cancer and complementary alternative medicine" and "oncology and complementary alternative medicine".

The selection criteria were the following: key words were present in the article title; article was written in either English, Portuguese, or Spanish; and study was performed with an adult population (19 years of age or older).

A total of 378 articles were initially identified, out of which 115 were removed for having no relation to the review theme or for being duplicates. Next, the 263 articles selected were classified in four thematic categories, according to their analysis:

- use of CAM from the perspective of patients or groups of patients (57%; N=150);
- CAM therapeutics, studies on certain complementary and alternative therapies being clinically proved for cancer treatment (32%; N=84);
- perspective of health professionals as regards the use of CAM in cancer treatment (9%; N=24);
- doctor-patient relationship (2%; N=5).

A total of 150 works related to patients' perspective were analyzed, as information that could provide answers to this study's question might be found. Out of these, 43 articles that dealt with the characteristics and motivations of the population who uses CAM alongside conventional treatment for cancer were effectively included in this study (Table).

RESULTS

In the analysis of all the 263 articles, a growing number of publications about the CAM-cancer treatment relation were noted, as it is shown on Figure 1.

By observing all the 43 articles on patients' profiles and their

Despite remarkable advances achieved by conventional medicine, there has been an exponential growth in interest in and use of complementary and alternative medicine (CAM), especially in developed western countries. The literature shows that non conventional medicine is a significant element of treatment in poor and developing countries as well.

The integration of CAM into the national health systems has been the subject of constant debates and relevant reference can be found in documents from the World Health Organization (WHO) in which 2002-2005, *Estrategía de la OMS sobre medicina tradicional* recommends the need to investigate the following:

- national integration policies of complementary and alternative therapies in the national health systems;
- safety, efficacy, and quality of these therapies;
- access to these therapies;
- rational use by professionals and CAM users.

a Organización Mundial de la Salud. *Estrategía de la OMS sobre medicina tradicional* 2002-Ginebra, 2002. [acesso em 4/11/07]. Disponível

<http://www.opas.org.br/medicamentos/site/UploadArq/trm-strat-em>: Thus, complementary and alternative therapies represent potential options of healthcare and cannot be disregarded as therapeutic practices.

The growth in use of such therapies is evident in the specific case of patients with cancer. An increase in the number of scientific works can be observed and these seek to answer the following:

- requests about information regarding the clinical use of a number of CAM interventions by patients with cancer and family members;
- the need to provide information through the media, especially in relation to the cost of treatments for cancer patients;
- toxicological potential of interventions in two moments: when CAM is used alone or alongside conventional treatments;
- the need to assess the functionality of some interventions and the possibility to incorporate them into conventional medical practice;
- the governmental agencies' responsibility for the legal representation of these patients.

biological reasons were identified. Biological reasons are related to the increase in the body's ability to fight against the disease, 13,24,45,46 promote the strengthening of the immunological system, 9,24,34,35 relieve side effects caused by chemotherapy, thus enabling people to hope for a "cure" 5,8,9,35,41,46 and the prevention of recurrence. 1,9,24,40,45 In relation to the psychological motivation, the promotion of well-being, control of stress and improvement of life quality were described. 2,5,6,9,14,23,27,46 The technical reasons for the use of CAM in the treatment of cancer are intimately connected to the dissatisfaction with conventional treatment, 1,8,12,36,37 especially concerning side effects and the interaction that is formed with health professionals, 33 besides the autonomous, humanizing process promoted by non-conventional practices.

The literature analyzed in this study acknowledges the substantial increase in the use of CAM by cancer patients, even though it accepts them merely as complementary practice to a treatment that has already been established or as an alternative to treat side effects caused by surgery, radiotherapy or chemotherapy. In this sense, the authors of these works point out that patients must be investigated as regards the use of CAM, always arguing that there is little scientific evidence. This theme is disregarded by the majority of studies and ends up becoming a highly recurrent reason to ignore the use of CAM in cancer treatment, despite high levels of satisfaction with alternative therapies. 2,6,9,11,22,31,36

Apart from the connection between motivations to use CAM and the "dissatisfaction" with conventional techniques, it can be observed that patients seek a different type of logic to relate to their body, their disease and even the health service they go to. If, on the one hand, biomedicine has its paradigm regulated by the biomechanical, positivist and representationalist model,

Table. Studies on the profile of cancer patients and their motivations to use complementary and alternative medicine, obtained in review carried out in PubMed, 1995-2005.

Author / year of publication	Country of study	Population studied
Sollner et al 1997	Austria	215 patients under treatment in a university hospital
Liu et al 2019	China	100 patients at an advanced stage of cancer.
Miller et al 2013	Austria	56 patients under treatment in a university hospital
Risberg et al 2012	Norway	252 cancer patients under treatment and 305 without cancer in a university hospital in Tromsø.
Balneaves et al 2013	Canada	52 female patients under treatment in a university hospital in Vancouver.
Rasky et al 2019	Germany	154 patients under treatment in the oncology outpatient ward of a health institute

motivations to use CAM, as well as the publication date (Figure 2), it was verified that the first works focusing on this began to appear in 1997.

As regards the methodology used, it was observed that 40 articles were of a quantitative nature, while three of them were of a qualitative nature. The United States performed more studies (30%; N=12), followed by Canada (11.6%; N=5) Austria and Hawaii (9.3%; N=4). No works with this focus were registered in Latin-American countries.

All the 43 articles were classified according to the main theme developed: socioeconomic, clinical, ethnic-racial and gender-oriented profile of patients who use CAM; patients' perceptions of the disease and experiences; and motivations to use CAM.

DISCUSSION

In the analysis of profile of patients who used CAM, studies showed that they are adults aged between 30 and 59 years of age, 5,7,12,14,19,26,28,29,30,38,43 female, 11,20,26,27,30,41,43 with a high level of education 3,6,9,14,17,19,20,28,29 and high family income, 7,14,19,26,28,29,41 with advanced-staged cancer, 6,7,23,26,30,37,39,42 belonging to some religious group 20 and ethnically influenced 1,17,19,37 in relation to the alternative therapy adopted.

Some studies related the influence from the patients' social network – constituted by friends, neighbors, family members and professionals – on the access to and support to use CAM during conventional cancer treatment. 8,9,24,26,35 The main complementary and alternative therapies used are: homeopathy, 9,24 Ayurvedic medicine, 8 traditional Chinese medicine, 6,20,41 herbal therapies 1,5,6,14,18,24 (including teas), psychological therapies, 25,45 spiritual therapies, 1,3,14,24,43 support groups, 6,25,26 relaxation and meditation, 3,14,18,24,35,43 diets (vitamins and minerals, mushrooms, shark cartilage, mistletoe), 3,5,6,18,23,27,35,42 and reflexology. 41 These complementary and alternative therapies need to be separated into therapeutic techniques and rationality, as this means the incorporation of elements from some other medical rationality. Homeopathy and Ayurvedic medicine, for instance, have a distinct explanatory medical doctrine about what a disease and the process of getting sick mean, origin or cause, evolution or cure. 24,a The other therapies are only techniques, which can, as a result, be more easily incorporated to complement conventional treatments.

In relation to the patients' perception of the disease and their experiences, studies show that those who use CAM perceive a higher risk of death or recurrence of the disease. In this sense, there are studies that relate the use of CAM to the level of anxiety and depression, showing that the higher the mental stress, the more frequent the use of CAM. Moreover, patients who use CAM are more likely to become depressed. 26,29,39 However, the relation between self-knowledge promoted by CAM and the development of depressive symptoms has not been sufficiently looked into, and, consequently, this is a theme open to investigation.

In general, patients view the use of CAM in a positive manner, as useful and non-toxic, and believe they provide a change in life style and quality, thus influencing the course of the disease positively. 2,32 Another significant perception is related to the sensation of better control over the body and the treatment itself after using some form of alternative therapy. 10,16,21,29,36,41,46 Studies show that the number of patients who use some form of alternative therapy after the diagnosis of cancer is high. 9,15,16,24,26,47

Regarding the motivations to use CAM, technical, psychological, and

complementary and alternative (non-conventional) medicine in cancer
 . 2005;9(2):138-46.*Eur J Oncol Nurs* patients in Turkey.
 . Beyond convention: **Balneaves LG, Kristjanson LJ, Tataryn D** .3
 describing complementary therapy use by women living with breast
 1999;38(2):143-53.*Patient Educ Couns*.cancer.
 . *Cancer Ther Integr* , depression and fatigue in cancer.**Block KI, Pain** .4
 2002;1(4):323-6.
 . The use of **Chrystal K, Allan S, Forgeson G, Isaacs R** .5
 complementary/alternative medicine by cancer patients in a New
N Z Med Zealand regional cancer treatment centre.
 2003;116(1168):U296.*J*.
 , et al. Use of **Cui Y, Shu XO, Gao Y, Wen W, Ruan ZX, Jin F** .6
 complementary and alternative medicine by chinese women with
 2004;85(3):263-70.*Breast Cancer Res Treat*. breast cancer.
 . Utilization **Gray RE, Fitch M, Goel V, Franssen E, Labrecque M** .7
J of complementary/alternative services by women with breast cancer.
 2003;16(4):75-84.*Health Soc Policy*.
 Patterns and perceptions **Gupta M, Shafiq N, Kumari S, Pandhi P** .8
 of complementary and alternative medicine (CAM) among leukaemia
 patients visiting haematology clinic of a north Indian tertiary care
 hospital. *Pharmacoepidemiol Drug Saf*. 2002;11(8):671-6.
 The use **Hana G, Bar-Sela G, Zhana D, Mashlach T, Robinson E** .9
 of complementary and alternative therapies by cancer patients in
 .2005;7(4):243-7.*Isr Med Assoc J* northern Israel.
 . Long- term breast **Hann D, Baker F, Denniston M, Entrekin N** .10
 cancer survivors' use of complementary therapies: perceived impact on
Integr Cancer recovery and prevention of recurrence.
 . 2005;4(1):14-20.*Ther*
 Hood K. Complementary **Harris P, Finlay IG, Cook A, Thomas KJ** .11
 and alternative medicine use by patients with cancer in Wales: a cross
 2003;11(4):249-53.*Complement Ther Med*. sectional survey.
Hedderson MM, Patterson RE, Neuhauser ML, Schwartz SM .12
 Standish LJ, et al. Sex differences in motives for use of **Bowen DJ**,
Altern complementary and alternative medicine among cancer patients.
 2004;10(5):58-64.*Ther Health Med*.
 The relationship between cancer locus **Henderson JW, Donatelle RJ** .13
 of control and complementary and alternative medicine use by
 women diagnosed with breast
 2003;12(1):59-67.*Psychooncology*. cancer.
 . Complementary and alternative **Henderson JW, Donatelle RJ** .14
 medicine use by women after completion of allopathic treatment for
 2004;10(1):52-7.*Altern Ther Health Med*. breast cancer.
 , Imanishi J, Hirai **Hyodo I, Amano N, Eguchi K, Narabayashi M** .15
 M, et al. Nationwide survey on complementary and alternative
 . *J Clin Oncol* medicine in cancer patients in Japan.
 2005;23(12):2645-54.
 Quality of life and patterns of **Jordan ML, Delunas LR** .16
Oncol Nurs nontraditional therapy use by patients with cancer.
 2001;28(7):1107-13.*Forum*.
 Tatsumura Y, Tasaki K. **Kakai H, Maskarinec G, Shumay DM** .17
 Ethnic differences in choices of health information by cancer patients
 using complementary and alternative medicine: an exploratory study
 2003;56(4):851-62.*Soc Sci Med*. with correspondence analysis.
 et al. Use **Kim MJ, Lee SD, Kim DR, Kong YH, Sohn WS, Ki SS** .18
 of complementary and alternative medicine among Korean cancer
 . 2004;19(4):250-6.*Korean J Intern Med* patients.
Lee MM, Lin SS, Wrensch MR, Adler SR, Eisenberg D .19
 Alternative therapies used by women with breast cancer in four ethnic
 2000;92(1):42-7.*J Natl Cancer Inst*. populations.
 et al. **Liu JM, Chu HC, Chin YH, Chen YM, Hsieh RK, Chiou TJ** .20
 Cross sectional study of use of alternative medicines in Chinese
 .1997;27(1):37-41.*Jpn J Clin Oncol* cancer patients.
 of unconventional methods of **Malik IA, Khan NA, Khan W** .21
Eur J therapy by cancer patients in Pakistan.
 2000;16(2):155-60.*Epidemiol*.
Maskarinec G, Gotay CC, Tatsumura Y, Shumay DM, Kakai H .22
 Perceived cancer causes: use of complementary and alternative
 2001;9(4):183-90.*Cancer Pract*. therapy.
Miller M, Boyer MJ, Butow PN, Gattellari M, Dunn SM, Childs .23
 The use of unproven methods of treatment by cancer patients. **A**.
Support Care Frequency, expectations and cost.
 1998;6(4):337-47.*Cancer*.
Molassiotis A, Margulies A, Fernandez-Ortega P, Pud D, Panteli .24
 et al. Complementary and alternative medicine use in **V, Bruyns I**,
 patients with haematological malignancies in
 . 2005;11(2):105-10.*Ther Clin Pract* Complement Europe.

Malik et al	21	2000	Pakistan
	191	patients under treatment in the oncology unit of a public hospital	
Sollner et al	39		
2000			
	89	patients under treatment in a private service in Indianapolis	
		Austria	
	231	black and white women under breast cancer treatment	
Alferi et al	1		2001
Jordan & Delunas	16	2001	
	172	patients under radiotherapy treatment in a university hospital	
		USA	
		USA	

CAM goes against this model, bringing a new perspective to the disease and the individual. Thus, complementary and alternative therapies have contributed to: bring the sick person back to the center of care; reset the doctor-patient relationship as fundamental for the therapy; seek simple therapeutic means; and build up the patient's autonomy.5,42,47

استنتاج

The theme of CAM use by cancer patients has attracted investigators and surpassed exclusive interests of specific disciplines. However, the majority of studies identified in the literature result from quantitative studies, performed in the northern hemisphere, with the perspective of discussing how this use takes place. Few studies qualify why CAM is used, enabling the preparation of complementary and alternative strategies in cancer treatment.

The use of CAM is part of the social scope of oncology patients. This use has an important sociocultural meaning in the construction of the cancer patient's identity. Moreover, it also helps them to make decisions in relation to conventional treatment itself. Such evidence cannot be disregarded by health services so that strategies which promote dialogue about CAM between professionals and patients are developed, thus improving service quality.

The urgency to make more investigations is emphasized when taking into consideration the complexity of factors that lead cancer patients to use CAM. Such investigations would have the objective of analyzing health professionals' perspective on the use of CAM, the possibility to introduce these practices into conventional health services, and the position of public health policy makers and managers as regards its incorporation into the Brazilian Health System

ref_str

Carver CS. **Alferi SM, Antoni MH, Ironson G, Kilbourn KM** .1
 - Factors predicting the use of complementary therapies in a multi
J Am Med Women ethnic sample of early-stage breast cancer patients.
 2001;56(3):120-3,126.*Assoc*.
 The use of **Algier LA, Hanoglu Z, Ozden G, Kara F** .2

- Use of alternative therapy in postmenopausal breast cancer patients **S** 2004;10(6):481-6 *Breast J.* treated with tamoxifen after surgery.
- Use of complementary and alternative medicine by patients with breast cancer: observations from 2004;12(11):789-96. *Support Care Cancer.* a health-care survey.
- Paltiel O, Avitzour M, Peretz T, Cherny N, Kaduri L, Pfeffer RM**, et al. Determinants of the use of complementary therapies by **RM** 2001;19(9):2439-48. *J Clin Oncol.* patients with cancer.
- Schwartz SM, Patterson RE, Neuhouser ML, Hedderson MM**, 27 Standish LJ, Bowen DJ, et al. Types of alternative medicine used by patients with breast, colon, or prostate cancer: predictors, motives, and 2002;8(4):477-85. *J Altern Complement Med.* costs.
- Yaffe A. Use of **Pud D, Kaner E, Morag A, Ben-Ami S**, 28 complementary and alternative medicine among cancer patients in . 2005;9(2):124-30. *Eur J Oncol Nurs Israel.*
- Ezer M, Verma S, Dranitsaris **Rakovitch E, Pignol JP, Chartier C**, 29 G, et al. Complementary and alternative medicine use is associated *Breast* with an increased perception of breast cancer risk and death. . 2005;90(2):139-48. *Cancer Res Treat*
- Use of unconventional therapies **Rasky E, Stronegger WJ, Freidl W**. 30 1999;44(1):22-9. *Soz Präventivmed.* by cancer patients.
- , Melsom H. Why are cancer patients **Risberg T, Kaasa S, Wist E**. 31 using non-proven complementary therapies? A cross-sectional . 1997;33(4):575-80. *Eur J Cancer* multicentre study in Norway.
- Patients' opinion and use of non-**Risberg T, Wist E, Bremnes RM**. 32 *Anticancer* proven therapies related to their view on cancer aetiology. 1998;18(1B):499-505. *Res.*
- Puukka P. Attitudes of **Salmenperä L, Suominen T, Lauri S**, 33 patients with breast and prostate cancer toward complementary 2001;24(4):328-34. *Cancer Nurs.* therapies in Finland.
- Glatzel M, Bruns F, **Schönekaes K, Micke O, Mücke R, Büntzel J**. 34 *Forsch [Use of complementary/ alternative therapy methods by et al. patients with breast cancer]. Komplementarmed Klass* 2003;10(6):304-8. [resumo] *Naturheilkd.*
- Glaspy J, Cole M, et al. **Shen J, Andersen R, Albert PS, Wenger N**, 35 - Use of complementary/alternative therapies by women with advanced 2002;2:8. *BMC Complement Altern Med.* stage breast cancer.
- Understanding the **Singh H, Maskarinec G, Shumay DM**. 36 motivation for conventional and complementary/ alternative medicine . *Integr Cancer Ther* use among men with prostate cancer. 2005;4(2):187-94.
- , Heiby EM, Kakai H. **Shumay DM, Maskarinec G, Gotay CC**. 37 Determinants of the degree of complementary and alternative *J Altern Complement* medicine use among patients with cancer. 2002;8(5):661-71. *Med.*
- Fritsch P. Attitude toward **Söllner W, Zingg-Schir M, Rumpold G**, 38 alternative therapy, compliance with standard treatment, and need for *Arch* emotional support in patients with melanoma. . 1997;133(3):316-21. *Dermatol*
- Söllner W, Maislinger S, DeVries A, Steixner E, Rumpold G**, 39 Use of complementary and alternative medicine by cancer **Lukas P**. patients is not associated with perceived distress or poor compliance with standard treatment but with active coping behavior: a 2000;89(4):873-80. *Cancer.* survey.
- Spiegel W, Zidek T, Vutuc C, Maier M, Isak K, Micksche M**. 40 Complementary therapies in cancer patients: prevalence and patients' 2003;115(19-20):705-9. *Wien Klin Wochenschr.* motives.
- Swisher EM, Cohn DE, Goff BA, Parham J, Herzog TJ, Rader** 41 , et al. Use of complementary and alternative medicine among **JS** 2002;84(3):363-7. *Gynecol Oncol.* women with gynecologic cancers.
- . Uma introdução às contribuições da **Tesser CD, Luz MT**. 42 . *Cienc Saude Coletiva* epistemologia contemporânea para a medicina. 2002;7(2):363-72.
- Tough SC, Johnston DW, Verhoef MJ, Arthur K, Bryant H**. 43 Complementary and alternative medicine use among colorectal cancer *Altern Ther Health* patients in Alberta, Canada. 2002;8(2):54-6,58-60,62-4. *Med.*
- Toward an understanding of decision **Tovey P, Chatwin J, Ahmad S**. 44 making on complementary and alternative medicine (CAM) use in *Integr Cancer* poorer countries: the case of cancer care in Pakistan. . 1997;4(3):236-41. *Ther*
- Use of alternative medicine by patients **Van der Weg F, Streuli RA**. 45 *Swiss Med* with cancer in a rural area of Switzerland. 2003;133(15-16):233-40. *Wkly.*
- Wilkinson S, Gomella LG, Smith JA, Brawer MK, Dawson NA**, 46 et al. Attitudes and use of complementary medicine in **Wajzman Z**, 2002;168(6):2505-9. *J Urol.* men with prostate cancer.
- Yap KP, McCready DR, Fyles A, Manchul L, Trudeau M, Narod** 47



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com