



International Journal Of Scientific And University Research Publication

ISSN No **301/704**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



GROWTH OF CULTURED WHITE LEG SHRIMP *LITOPENAEUS VANNAMEI* (BOONE, 1931) OF BRACKISH WATER CULTURE SYSTEM IN WINTER SEASON WITH ARTIFICIAL DIET

Danya Babu Ravuru || Department of Zoology and Aquaculture
Acharya Nagarjuna University
Guntur
Andhra Pradesh-522510
India.

The White leg shrimp *Litopenaeus vannamei* culture was conducted from three ponds each one of 0.5 ha for the study. Semi-Intensive culture system was selected under brackish water

sh water conditions. Stocking densities (post larvae) were taken from 3 samples; each one contains 50 Numbers/m². In winter season in month of November to February, the water quality parameters were measured fortnightly in a month at 7 a. m. The production was 3200, 3318 and 3459 kg and FCR was 1.43, 1.51 and 1.46 and the final growth was 16.50, 17.00 and 17.50 g /90, 92 and 94 days for P1, P2 and P3, respectively.

L. vannamei; Temperature, Salinity, Density; Feed; Growth and

مقدمة

13.5-16.5		13.0-16.0	Temperature (°C)
6.5-12.5	6.0-12.5	6.0-12.0	Salinity (ppt)
3.5-4.2	3.5-4.0	3.4-4.0	DO (ppm)

Table 1: Average water quality parameters

						Pond
	75	60	45	30	15	
	13.00	10.00	7.00	4.00	2.00	P1

Table 2: Fortnightly growth performance (g)

						Pond
	75	60	45	30	15	
	13.50	10.50	7.50	4.50	2.00	P2

Table 3: Fortnightly growth performance (g)

						Pond
94	75	60	45	30	15	
17.50	13.50	10.50	7.50	4.50	2.00	P2

Table 4: Fortnightly growth performance (g)

Pond 3	Pond 2	Pond 1	Details				
0.5	0.5	0.5	Area (ha)				
2,50,000	2,50,000	2,50,000	Initial stocking (Numbers)				
50	50	50	Density/m ²				
02/11/2012	02/11/2012	02/11/2012	Stocking Date				
PL	PL	PL	PL stocking (Days)				
06/02/2013	04/02/2013	02/02/2013	Harvest Date				
17.50	17.00	16.50	Harvest size (gm)	P3 range	P2 range	P1 range	Parameters
57.1	58.8	60.6	Count (numbers/kg)	7.5-8.2		7.0-7.5	PH

Because of the high demand for shrimps in Japan, the United States and Europe, shrimp aquaculture has expanded rapidly in all around the world, especially in tropical areas, such as Southeast Asia and , which *vannamei* Latin America [1]. Among all species of shrimp, *L. represent* over 90% of shrimp culture in the Western hemisphere, is the most commonly cultured shrimp in Central and South American countries, China, and Thailand [2-4]. India has the one of the longest line of 8118 km.

Material and Methods

at beginning the *vannamei* The shrimp 15 days old post larvae *L. study* collected from BMR hatchery. The winter season experiments survival were 82, 84 and 86%, respectively. *vannamei* the species *L. Water* depth maintained 8 ft. The water takes from creek pumped to 3 ponds by PVC pipe (size 2½ inch). The PH, temperature, salinity and DO ranges up to 7.0-8.2, 13.0- 16.5°C, 6.0-12.5 ppt and 3.4-4.2 ppm/day for P1, P2 and P3. The artificial diet was given made by *Manamei* feed (protein% 35 (Feed No. 1, 2, 3 and 3S) and 34(Feed No. 3M)).

Results

In the experiment the water quality parameters are presented (Table well grow, the body weight increased *L.vannamei* 1). The species 3.0-3.5 g and ADG 0.18 g (P1, P2 and P3)15 days in Indian climate conditions, which is better than other countries. In the culture system the growth rate increased due to the artificial feed supplementation in the season. The mean average weights of the shrimp at harvest were 16.5, 17.0 and 17.5 g (Tables 2, 3 and 4) and production was 3200, 3318 and 3459 kg and FCR was 1.36, 1.40 and 1.46/90, 92 and 94 for P1, P2 and P3, respectively.

Discussion

A large number of shrimp could be assembled on the aquaria bottom from the artificial substrates [5]. Many studies have illustrated that artificial substrates could increase shrimp growth and survival [6,7]. Abiotic factors such as temperature and salinity may also affect the protein requirement [8]. “Booster” mineral is given to 3 ponds depending on biomass for development the minerals. “Opti oxygen” controls the DO. “AQ lite” for bottom clears. The shrimp maintained at 35°C had the highest rate of food consumption Arenda et al. [9] recorded the average growth rate of 0.38 g/wk in the 90 shrimp/ m² and lowest in the 180 shrimp/m² (0.33 g/wk). My observation at 7-8.2°C the shrimp highest growth rate 4 g/fortnightly (50 shrimp/m²) and lowest growth 3.5 g/fortnightly (50 shrimp/m²) /94 and 92 days.

- [density. Aquaculture 293: 42-48.](#)
Protein and amino acids. World Aquaculture **Guillaume J (1997)**.8
Society, Baton Rouge, USA.
- [White shrimp *Penaeus* **Arenda M, Perez EP, Gasca-Leyva E \(2008\)**.9](#)
[culture in fresh water 3densities: condition state based on vannamei](#)
Aquaculture 283: 13-18. [length and weight.](#)
[Response**Cawthorne DE, Beard T, Davenport J, Wickins J \(1983\)**.10](#)
[Penaeus monodon Fabricius to natural and artificial sea of juvenile](#)
Aquaculture 32:165-174. [water of low salinity.](#)
[Lethal levels of low dissolved oxygen**Allen GL, Maguire GB \(1991\)**.11](#)
[short-term oxygen stress on subsequent growth of and effect of](#)
Aquaculture 94: 27-37. [juvenile *Penaeus monodon*.](#)
[Transport limitation of oxygen in shrimp**Garcia A, Brune DE \(1991\)**.12](#)
pond. Aqua Eng 10: 269-279. [culture](#)
Crustacean forming Black Well **Lee DOC, Wickins JE (1992)**.13
Scientific Publications. Oxford.
- [Parado-Estpa EED, Llobera A, Villaluz A, Saldes R \(1993\).14](#)
[metamorphosis of *Penaeus monodon* Larvae at differentSurvival and](#)
J Aqua 45: 3-7. [salinity levels. Israel](#)
[Growth of cultured white**Danya Babu R, Jagadish Naik M \(2014\)**.15](#)
[Litopenaeus vannamei \(Boone 1931\) of brackish water of shrimp](#)
[summer season with artificial diet. Adv Appl Sci culture system in](#)
[Res 5: 25-28.](#)

101	98	95	Doc	
86	84	82		Survival (%)
1.46	1.40	1.36		FCR
0.18	0.18	0.18		ADG (g)
3459	3318	3200		Production (kg)
2486.70	2359.40	2352		Total feed (kg)
Rs 20.50	Rs 20.50	Rs 20.50		Seed cost/kg shrimp

Table 5: Average Cost Analysis

Early morning Dissolved Oxygen concentration was between 3 and 5 mg/l-1; salinity was above 15‰ during the first week of grow out pond, which is preferable for post larvae [10-15]. In early morning I reported Dissolved Oxygen concentration was between 3.4-4.2 ppm and salinity was between 6-12.5 ppt. For each pond cost analysis was worked out. Production cost for 1 kg shrimp (60.6, 58.8 and 57.1 counts) was calculated as Rs 300, 350 and 380. The feed cost was Rs 71.84/kg, followed by seed cost Rs 20.50/kg. Totally feed was used 2352, 2359 and 2359 kg. According to Danya Babu Ravuru and Jagadish Naik Mude, in summer season better production (kg) and profit (Rs) was 8337, 8932 and 9450; 218.34, 243.63 and 272.66/P1, P2 and P3 for 120, 123 and 126 days and 3200, 3318 and 3459; 102, 152 and 182/ P1, P2 and P3 for 90, 92 and 94 days (Table 5).

استنتاج

In the present study, it has been observed, Temperature, Salinity, Dissolved Oxygen, Density and Survival. The culture is successful in brackish water and *L. vannamei* shrimp shrimp growth was increased with artificial feed.

Authors are thankful to the Farmer and **Acknowledgement** Owner of the culture ponds K. Ramana (Neeli Aqua Pvt Ltd) in Chinaganjam Village, Prakasakm District, for their encouragement and provided facilities up to harvest.

ref_str

- [Lombardi JV, De Almeida MHL, Toledo LPR, Salee BOJ, De Paula .1](#)
[Cage Polyculture of the Pacific white shrimp *Litopenaeus* EJ \(2006\)](#)
[Philippines Sea weed *Kappaphycus alvarezii*, *vannamei* and the](#)
[Aquaculture 258: 412-415.](#)
- [Acute**Frias-Espericueta MG, Voltolina D, Osuna-López JJ \(2001\)**.2](#)
[cadmium, mercury and lead to white leg shrimp toxicity of](#)
[larvae. Bulletin of Environmental \(Litopenaeus vannamei\) post](#)
[Contamination and Toxicology 67: 580-586.](#)
- [Mc Graw WJ, Davis DA, Teichert-Coddington D, Rouse DB .3](#)
[of *Litopenaeus vannamei* post larvae to low Acclimation\(2002\)](#)
[endpoint and rate of salinity salinity: influence of age, salinity,](#)
[reduction. J World Aquacul Soc 33: 78-84.](#)
- [Suitability studies of inland**Saoud IP, Davis DA, Rouse DB \(2003\)**.4](#)
[for *Litopenaeus vannamei* culture. Aquaculture 217: well waters](#)
[373-383.](#)
- [Effects of artificial**Zhang B, Li WH, Huang YJW, Xu RL \(2010\)**.5](#)
[on the growth, survival and spatial distribution of substrates](#)
[intensive culture condition. Iran J Fish SciLitopenaeus vannamei in the](#)
[9: 293-304.](#)
- [Effects of artificial substrate and stocking density on**Moss SM \(2004\)**.6](#)
[nursery production of Pacific White shrimp *Litopenaeus* the](#)
[Aquaculture soci 35: 537-542. *vannamei*. J World](#)
[-High**Arnold SJ, Coman FE, Jackson CJ, Groves SA \(2009\)**.7](#)
[water-exchange production of juvenile tiger shrimp, intensity, zero](#)
[evaluation of artificial substrates and stocking *Penaeus monodon*: An](#)



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com