



International Journal Of Scientific And University Research Publication

ISSN No **2364/2017**

Listed & Index with
ISSN Directory, Paris



Multi-Subject Journal



ACUTE TOXICITY OF ALPHA-CYPERMETHRIN TO OLIGOCHAETE WORM, BRANCHIURA SOWERBYI (BEDDARD, 1982) ALONG WITH THEIR BEHAVIOURAL RESPONSES.

Dr.Dip Mukherjee || Assistant Professor

Post Graduate Department of Zoology

Bidhannagar College

EB-2

Sector-1

Salt Lake City

Kolkata - 700064

India.

Acute toxicity of alpha-cypermethrin to *Branchiura sowerbyi* and their behavioural changes were evaluated in the present study. The 24, 48, 72 and 96 h LC₅₀ values of

alpha-cypermethrin to *B. sowerbyi* were 43.39, 39.75, 22.58 and 13.41 µg/l respectively. A significant variation ($p < 0.05$) between mortality rate of worm and concentrations of the test chemical at all the exposure times (24, 48, 72 and 96 h) was observed. The mortality rate also significantly varies ($p < 0.05$) with the exposure time at all the concentrations except 48h. In addition, a dose dependent change in the behavioural pattern of the worm was recorded. It was directly proportional to the increasing concentration of alpha-cypermethrin.

Acute toxicity, alpha-cypermethrin, *Branchiura sowerbyi*, behavioural

مقدمة

Initially, rough range finding tests were conducted for the test organism to determine the dose range at which mortality occurs (data not shown). The selected test concentrations of alpha-cypermethrin were finally used for the determination of LC₅₀ values for . Ten test organisms (mean length 18.5 ± 6.22 *Branchiura sowerbyi* the mm) were used in each replicate. The number of dead organisms was counted at every 24h during the experiment. The dead animals were removed immediately to avoid any organic decomposition. A constant amount of test medium was replaced every 24h by freshwater and the desired quantity of alpha-cypermethrin was immediately added to water to assure a fixed concentration of the toxicant in solution and also to avoid other interfering factors affecting the test animals' performance. Water chemical analysis and the bioassays were done following the methods outlined in American Public Health Association (2012). Similar technique was also followed by previous workers (Badanthadka & Mehendale, 2005; Mukherjee & Saha, 2012). The LC₅₀ values for 24, 48, 72 and 96h along with 95% confidence limits were estimated by a computer program (US EPA, 1999).

The behavioural changes of the exposed worms at each concentration of alpha-cypermethrin were also recorded during the bioassay. The data on percent mortality was subjected to analysis of variance using the computer software provided by the R Development Core Team (2011) followed by Duncan's multiple range test to determine significant differences among means at different times of exposure and concentrations (Gomez & Gomez, 1984).

Results and Discussion

Branchiura The lethal concentration of alpha-cypermethrin to was *B. sowerbyi* is summarized in Table 1. No mortality of *sowerbyi* recorded in control during the experiment. The mortality rate (%) of the test animals was significantly increased ($p < 0.05$) from control with increasing concentration of the test chemical except 0.8 µg/l at all the exposure times (24, 48, 72, 96h). On the other hand, the mortality rate also varied significantly ($p < 0.05$) with the progress of exposure time except 48h at all the concentrations (Table 2).

With the increasing concentration of test chemical and the progress showed different irregular behaviors *B. sowerbyi* of exposure time, which are shown in Table 3. The control worms were active throughout the test period and showed clumping tendency with normal movement. The clumping tendency of the exposed worms was gradually reduced with increasing concentrations and times of exposure. The worms showed rapid movement at all the higher doses.

Alpha-cypermethrin, an active isomer of cypermethrin, is used to control the common domestic insects and pests belonging to families like Lepidoptera and Coleoptera infesting vegetables, citrus fruits, cotton, rice, etc. (Yordanova et al., 2009). Cypermethrin has been reported to be used against lice invasion in fishery (Das & Mukherjee, 2003). Alpha-cypermethrin is a synthetic pyrethroid which acts as a contact and stomach poison. It is classified as a Schedule 6 poison in the Standard for the Uniform Scheduling of Drugs and Poisons (Sarikaya, 2009). Alpha-cypermethrin acts on pre-synaptic neurons of organisms where it prevents the transmission of nerve impulses through the axon by blocking the passage of sodium ions across the channels in nerve fibers (Yordanova et al., 2009). It is non-systemic in nature and is moderate to highly toxic to diverse aquatic and terrestrial life (Mokry & Hoagland, 1990; Chamber, 1994). Very little information is available on the acute toxicity of alpha-cypermethrin on aquatic invertebrates (Yordanova et al., 2009). It is highly toxic to fish. The 24 h LC₅₀ values of alpha cypermethrin to silver barb and common mirror were 20.0 and 4.50 µg/l respectively (Grayson et al., 1990). There is no specific report on the lethality of alpha-cypermethrin on aquatic annelids particularly on the benthic ones that play a vital role in detritus food chains. This sprouted the present need to investigate the acute toxic effects of and their behavioural *Branchiura sowerbyi* alpha-cypermethrin to responses

Materials and Methods

Test organism used in the bioassay was the benthic oligochaete (Class: Oligochaeta, Family: Tubificidae). *Branchiura sowerbyi* worm. The test organisms were collected from local non-polluted sources. The worms were allowed to acclimatize in the test water for 3 days before the experiment.

Commercial grade alpha-cypermethrin (10% w/w, EC) was collected from the local market. Static replacement bioassays with the was conducted in 500 ml Borosil glass beakers each *sowerbyi* containing 250 ml unchlorinated tap water (Temperature 28.3 ± 0.4 °C, pH 7.1 ± 0.2 , Free CO₂ 9.9 ± 0.5 mg/l, Dissolved Oxygen 5.5 ± 0.3 mg/l, Total alkalinity 181 ± 8.6 mg/l as CaCO₃, Hardness 119 ± 6.2 mg/l as CaCO₃). A set of four beakers were exposed to each concentration of alpha-cypermethrin. Each set of tests was accompanied by four replicates of control.

Stock solution of the test chemical and its dilutions were made following the method of American Public Health Association (2012).

00am ± 0.43	00am ± 0.43	0.8
10bm ± 0.00	10bm ± 0.43	8
20cm ± 0.71	20cm ± 0.71	20
50dm ± 0.50	40dm ± 0.87	40
60emn ± 0.83	50em ± 0.50	60
70fm ± 0.00	70fm ± 0.43	80
80gm ± 0.00	80gm ± 0.50	100
90hm ± 0.00	90hm ± 0.43	120
100im ± 0.00	100hm ± 0.43	150

Table 3: Impact of alpha-cypermethrin on behavioural responses of *Branchiura sowerbyi* (M: movement; CT: clumping tendency; MS: mucus secretion; WE: wrinkling effect; -: none; +: mild; ++: moderate; +++: strong) exposed to various concentrations during different hours of exposure

		Concentrations (µg/l)					Time of exposure
		100	80	60	40	20	8
+++	+++	++	++	+	-	M	24h
	-	-	++	++	+++	+++	CT
	++	++	+	-	-	-	MS
	+++	+++	+	+	-	-	WE
+++	+++	++	++	+	-	M	48h
	-	-	+	++	+++	+++	CT
	++	++	+	+	-	-	MS
	+++	+++	++	+	-	-	WE
+++	+++	++	++	-	-	M	72h
	-	+	-	++	+++	+++	CT
	++	++	+	-	-	-	MS
	+++	+++	++	+	-	-	WE

The rate of movement was increased gradually with the increasing concentrations and time of exposures. The wrinkling effect and gradually increased at 60, 80, and *B. sowerbyi* mucous secretion of 100 µg/l of alpha-cypermethrin at all the exposure times.

B. In the present study, the 96h LC50 value of alpha-cypermethrin to *B. sowerbyi* is 13.41 µg/l. The aquatic invertebrates and fish are most vulnerable to alpha-cypermethrin as its rate of metabolism and elimination are much slower in their body (WHO, 1992; Greulich & Pflugmacher, 2003). The 96h LC50 value of cypermethrin *Tilapia nilotica*, *Cyprinus carpio*, *Salmo trutta*, *Salmo gairdneri* for were 2.20, 0.90-1.10, 1.20, 0.50 and *Scardinius erythrophthalmus* and 0.40 µg/l respectively (Sarikaya, 2009).

The present investigation indicates that the 96h LC50 value of alpha is slightly higher than the other aquatic *B. sowerbyi* cypermethrin to invertebrates and vertebrates. The 48h EC50 value of alpha *Gammarus* was 0.8 µg/l and for *Daphnia magna* cypermethrin for 24h LC50 value was 0.3 µg/l (Yordanova et al., 2009). Yilmaz et al. (2004) and Sarikaya (2009) reported that the 96h LC50 value of alpha-cypermethrin for guppy and Nile tilapia were 9.43 µg/l and 5.99 µg/l respectively.

Branchiura sowerbyi The LC50 value of alpha-cypermethrin to provide useful data in the determination of safe level for the agricultural effluents before their release to the natural water resources. The evaluation of the toxicity of alpha-cypermethrin is not sufficient for assuring safe level of the receiving water bodies. In addition, potential risk from alpha-cypermethrin metabolites should also be considered to get a more accurate picture in terms of toxicity.

Table 1: Median lethal concentration (LCT) of alpha-cypermethrin to the *Branchiura sowerbyi* at different hours of exposure with 95% confidence limits

Concentration (µg/l)	48h	72h
43.39	39.75	24h
(28.55-59.14)	(25.62-54.48)	<i>Branchiura sowerbyi</i>

Table 2: Mean values (± SD) of mortality rate (%) of *Branchiura sowerbyi* exposed to different concentrations of alpha-cypermethrin at different hours of exposure (24h, 48h, 72h, 96h). Mean values within columns indicated by different superscript letters (a-h) and mean values within rows indicated by different superscript letters (m-o) are significantly different

Hours of exposure	48	72
Concentration (µg/l)	24	24
0.0	0.0	0.0
0.0	0.0	0.0
0.0	0.0	0.0

+++	+++	+++	++	+	-	M	96h
-	-	-	++	++	++	CT	
+++	+++	++	++	+	-	MS	
+++	+++	++	+	+	-	WE	

استنتاج

We are thankful to the Head, Postgraduate Department of Zoology and the Principal of Bidhannagar College for extending all kinds of help to carry out the present work.

ref_str

- (APHA) (2012). Standard **American Public Health Association**. 1 methods for the examination of water and wastewater. (Eds. Rice, E.W., Baird, R.B., Eaton, A.D. & Clesceri, L.S.) American Public Health Association, American Water Works Association, Water Environment Federation, Washington
- . (2005). Chlorophenols. **DC. Badanthadka, M. & Mehendale, M.H.** 2 pp. 567-568. In: Encyclopedia of Toxicology-1. (Ed. Elsevier. Chambers, J. (1994). Toxicity **Wexler, P.) Academic Press**, 3 of pesticides. In: Basic Environmental Toxicity (Eds. Cocherhan, L.G. & Shane, B.S.) CRC Press, Boca Raton, 185-198.
- (2003). Toxicity of cypermethrin in **Das, B.K. & Mukherjee, S.C.** 4 Labeo rohita fingerlings: biochemical, enzymatic and haematological consequences. Comparative Biochemistry & Physiology, 134(C), 109-121.
- . & Copping, L.G. (1990). Pest **Grayson, B.T., Green, M.B.** 5 Management in Rice Conference, London, UK, pp. 455-464.
- . (1984). Statistical procedures for **Gomez, K.A. & Gomez, A.A.** 6 agricultural research, 2nd Ed. New York, Wiley.
- (2003). Differences in susceptibility **Greulich, K. & Pflugmacher, S.** 7 of various life stages of amphibians to pesticide exposure. Aquatic Toxicology, 65, 329-336.
- . (1990). Acute toxicities of five **Mokry, L.E. & Hoagland, K.D.** 8 synthetic pyrethroid insecticides to Daphnia magna and Ceriodaphnia dubia. Environmental Toxicology and Chemistry, 9(8), 1045-1051.
- (2012). Acute toxicity of 2, 4, **Mukherjee, D. & Saha, N.C.** 9 6-Trichlorophenol to copepod, Cyclops viridis and oligochaete worm, Branchiura sowerbyi. Environment and Ecology, 30(3C) 1165-1170.
- (2011). R: A language and environment **R Development Core Team**. 10 for statistical computing. R Foundation for Statistical Computing, <http://www.Rproject.org/> Vienna, Austria. ISBN 3-900051-07-0, URL Accessed 8 Oct 2012.
- (2009). Investigation of Acute Toxicity of Alpha- **Sarikaya, R.** 11 ypermethrin on Adult Nile Tilapia (Oreochromis niloticus **Turkish Journal of Fisheries and Aquatic Sciences**, 9, 85-89. L.). **US**
- Probit program version 1.5. Ecological Monitoring **EPA. (1999).** 12 Laboratory, US Research Division, Environmental Monitoring Systems Environmental Protection Agency, Cincinnati, Ohio
- . Accessed 14 Jan 2012. <http://www.epa.gov/nerleerd/stat2.htm> 45268.
- , (1992). Alpha- cypermethrin. **World Health Organization**. 13 Environmental Health Criteria. World Health Organization, Geneva.
- & Erbasli, K. (2004). Acute toxicity of alpha- **Yilmaz, M., Gul, A.** 14 ypermethrin to guppy (Poecilia reticulata, Pallas, 1859). Chemosphere, 56, 381-385.
- Stoyanova, T., Traykov, I. & Boyanovsky, B. (2009). **Yordanova, V.,** 15 Toxicological Effects of Fastac Insecticide (Alpha-cypermethrin) to Daphnia magna and Gammarus pulex. Biotechnology & Biotechnological Equipment, 23(2), Special edition, 393-395.



IJSURP Publishing Academy

International Journal Of Scientific And University Research Publication
Multi-Subject Journal

Editor.

International Journal Of Scientific And University Research Publication



+965 99549511



+90 5374545296



+961 03236496



+44 (0)203 197 6676

www.ijsurp.com